

REMARKS ON NAVIER-STOKES EQUATIONS WITH MEASURES AS DATA

B. MICHAUX

Laboratoire de Modélisation et Calcul—IMAG, Université Joseph Fourier
 BP 53X, 38041 Grenoble Cédex, France

J. M. RAKOTOSON

Département de Mathématiques, Faculté des Sciences de Poitiers
 40 Avenue du Recteur Pineau, 86022 Poitiers Cédex, France

(Received and accepted May 1993)

Abstract—We show the existence of a solution to the Navier-Stokes equation taking the vorticity ω as the unknown: $\omega_t + A\omega + B\omega = \mu$, $\omega(0) = \omega_0$. Here, ω_0 and μ are bounded Radon measures. This study is motivated by a numerical approximation which will be given in a forthcoming work [1].

1. INTRODUCTION

Usually, in the Navier-Stokes equations for an incompressible fluid, the unknowns are the velocity V and the pressure p (see [2]):

$$(P_1) \begin{cases} \frac{\partial V}{\partial t} - \nu \Delta V + (V \cdot \nabla)V + \text{grad} p = f \text{ in } Q = (0, T) \times \Omega, \\ \text{div} V = 0 \text{ in } Q. \end{cases}$$

Assume $\Omega \subset \mathbb{R}^2$, physicists also consider as unknowns the vorticity $\omega = (\text{curl } V)\vec{k}$ and the stream function ψ satisfying $V = \text{curl}(\psi\vec{k})$, where \vec{k} is the unit vector orthogonal to the plane \mathbb{R}^2 . If we take the “curl” of the first equation and take the projection on the k -axis, we get from (P_1) :

$$(P_2) \begin{cases} \frac{\partial \omega}{\partial t} - \nu \Delta \omega + (V \cdot \nabla)\omega = \mu \text{ in } Q, \\ -\Delta \psi = \omega \text{ in } Q. \end{cases}$$

We supplement (P_2) with the additional conditions that $\omega = \psi = 0$ on $(0, T) \times \partial\Omega$ and $\omega(0, \cdot) = \omega_0$; ω_0 as well as μ are bounded Radon measures on Ω . If $\Omega = \mathbb{R}^2$, this problem was already considered by G.H. Cottet [3,4] when $\mu = 0$. So, we will assume that Ω is a bounded domain. Although the boundary condition on ω is not the usual one it is here a relevant one as our study is above all motivated by the numerical approximation of (P_2) in a large domain Ω .

2. NOTATIONS AND MAIN RESULTS

We denote by $M(Q)$ (resp., $M(\Omega)$) the set of bounded Radon measures on Q (resp., on Ω), and we set for $m \geq 1$ and $p \in [1, +\infty]$:

$$Z^{m,p}(\Omega) = \bigcap_{s \in [1,p[} W^{m,s}(\Omega), \quad Z_0^{m,p}(\Omega) = Z^{m,p}(\Omega) \cap W_0^{1,1}(\Omega).$$

Now, we define the B -functions on $Z_0^{1,4/3}(\Omega)$ by first introducing the map

$$L : Z_0^{1,4/3}(\Omega) \mapsto Z_0^{2,4}(\Omega) \text{ with } L\omega = \psi \text{ if and only if } -\Delta \psi = \omega \text{ in } \Omega, \psi = 0 \text{ on } \partial\Omega.$$

We set $B(\omega, \varphi) = \frac{\partial}{\partial x_2} L\omega \frac{\partial \varphi}{\partial x_1} - \frac{\partial}{\partial x_1} L\omega \frac{\partial \varphi}{\partial x_2}$ for (ω, φ) in $Z_0^{1,4/3}(\Omega) \times Z_0^{1,4/3}(\Omega)$ and $B\omega = B(\omega, \omega)$.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

As a property of B , we have the following.

LEMMA 1. Let Φ be a bounded lipschitz function from \mathbb{R} into \mathbb{R} , then for all $\omega \in Z_0^{1,4/3}(\Omega)$, we have

- (i) $B\omega \in L^p(\Omega)$, $1 \leq p < \frac{4}{3}$;
- (ii) $\int_{\Omega} B\omega(x)\Phi(\omega(x)) dx = 0$;
- (iii) $\forall \varphi \in W_0^{1,+\infty}(\Omega)$, $\int_{\Omega} B\omega\varphi dx = - \int_{\Omega} \omega B(\omega, \varphi) dx$.

PROOF. For $\omega \in Z_0^{1,4/3}(\Omega)$, $\text{grad}(L\omega) \in L^\infty(\Omega)^2$; so we have (i). We write

$$\int_{\Omega} B\omega\Phi(\omega) dx = \int_{\Omega} \frac{\partial}{\partial x_2} L\omega \frac{\partial}{\partial x_1} \left(\int_0^\omega \Phi(\sigma) d\sigma \right) dx - \int_{\Omega} \frac{\partial}{\partial x_1} L\omega \frac{\partial}{\partial x_2} \left(\int_0^\omega \Phi(\sigma) d\sigma \right) dx.$$

By integration by parts, we get (ii), and (iii) is also derived by an integration by parts.

The problem (P_2) is equivalent to solving the problem $\omega_t + A\omega + B\omega = \mu$ on Q , $\omega(t) \in Z_0^{1,4/3}(\Omega)$ for a.e., $t \in [0, T]$ where $A\omega = -\nu\Delta\omega$, $\nu > 0$.

THEOREM 1. (Existence of a weak solution) Let $\mu \in M(Q)$ and $\omega_0 \in M(\Omega)$, then there exists $\omega \in \bigcap_{q \in [1, \frac{4}{3}[} L^q(0, T; W_0^{1,q}(\Omega))$ satisfying

$$- \int_Q \omega \varphi_t dx dt - \int_{\Omega} \varphi(0, x) d\omega_0 + \nu \int_Q \nabla \omega \nabla \varphi dx dt - \int_Q \omega B(\omega, \varphi) dx dt = \int_Q \varphi(t, x) d\mu,$$

for all $\varphi \in C_0^\infty(\mathbb{R}^3)$, which is zero in a neighborhood of $(0, T) \times \partial\Omega$ and $\{T\} \times \Omega$.

REMARK. We prove below that $\omega B(\omega, \varphi)$ is in $L^1(Q)$.

3. PROOF OF THEOREM 1

We follow the same scheme as in [5]. First of all, we consider μ^ε in $C_0^\infty(Q)$ (resp., ω_0^ε in $C_0^\infty(\Omega)$) with μ^ε (resp., ω_0^ε) remaining in a bounded set of $L^1(Q)$ (resp., in $L^1(\Omega)$), ($\varepsilon > 0$). Using the result in R. Temam's book [2], we derive the existence of a function $\omega^\varepsilon \in L^2(0, T; H^2(\Omega) \cap H_0^1(\Omega))$ satisfying: (P_ε) : $\omega_t^\varepsilon + A\omega^\varepsilon + B\omega^\varepsilon = \mu^\varepsilon$ in $\mathcal{D}'(Q)$, $\omega^\varepsilon(0) = \omega_0^\varepsilon$.

From Lemma 1 and integration by parts, we derive the following lemma.

LEMMA 2. Let Φ be a bounded lipschitz function from \mathbb{R} into \mathbb{R} , then for all $t \in [0, T]$:

$$\begin{aligned} \int_{\Omega} dx \int_0^{\omega^\varepsilon(t,x)} \Phi(\sigma) d\sigma - \int_{\Omega} dx \int_0^{\omega_0^\varepsilon(x)} \Phi(\sigma) d\sigma + \nu \int_0^t \int_{\Omega} \nabla \omega^\varepsilon \nabla \Phi(\omega^\varepsilon) dx d\sigma \\ = \int_0^t \int_{\Omega} \Phi(\omega^\varepsilon) \mu^\varepsilon dx d\sigma. \end{aligned}$$

Choosing various functions Φ , as in [5], in Lemma 2 and applying Lemma 1 of [5], we deduce the following lemma.

LEMMA 3. ω^ε remains in a bounded set of $L^q(0, T; W_0^{1,q}(\Omega))$ for all $q \in [1, \frac{4}{3}[$ and also in $L^\infty(0, T; L^1(\Omega))$ as $\varepsilon \rightarrow 0$.

LEMMA 4. There exists $r > 1$ such that $\omega^\varepsilon \text{grad}(L\omega^\varepsilon)$ remains in a bounded set of $L^r(\Omega)^2$.

PROOF OF LEMMA 4. For a fixed $q \in [1, \frac{4}{3}[$, we set $q^* = 2q/2 - q$. Consider $\eta > 0$ such that $q^*\eta < (q-1)(q^*-1)(1+\eta)$. By interpolation and Lemma 3, we have

$$|\omega^\varepsilon(t)|_{L^{1+\eta}(\Omega)} \leq |\omega^\varepsilon(t)|_{L^1(\Omega)}^\theta |\omega^\varepsilon(t)|_{L^{q^*}(\Omega)}^{1-\theta} \leq c |\omega^\varepsilon(t)|_{L^{q^*}(\Omega)}^{1-\theta}, \quad \forall t > 0.$$

Note that $1 - \theta = q^* \eta / (1 + \eta)(q^* - 1)$. From the definition of $L\omega^\varepsilon$, Calderon-Zygmund's theorem and the preceding estimate, we get:

$$\begin{aligned} |\omega^\varepsilon(t) \operatorname{grad}(L\omega^\varepsilon)(t)|_{L^{1+\eta}(\Omega)} &\leq |\operatorname{grad}(L\omega^\varepsilon)(t)|_{L^\infty(\Omega)} |\omega^\varepsilon(t)|_{L^{1+\eta}(\Omega)} \\ &\leq c |\omega^\varepsilon(t)|_{L^{q^*}(\Omega)} |\omega^\varepsilon(t)|_{L^{q^*}(\Omega)}^{1-\theta} = c |\omega^\varepsilon(t)|_{L^{q^*}(\Omega)}^{2-\theta}. \end{aligned}$$

Since $q/(2 - \theta) > 1$, we have for $r = \inf(q/2 - \theta, 1 + \eta)$,

$$|\omega^\varepsilon \operatorname{grad}(L\omega^\varepsilon)|_{L^r(Q)} \leq c |\omega^\varepsilon|_{L^q(0,T;W_0^{1,q}(\Omega))}^{2-\theta} \leq \text{constant}.$$

COROLLARY OF LEMMAS 3 AND 4.

- (i) There exists $s > 1$ such that: ω_t^ε remains in a bounded set of $L^1(0, T; H^{-s}(\Omega))$.
- (ii) For all φ in $C^\infty(\overline{Q})$, $\omega^\varepsilon B(\omega^\varepsilon, \varphi)$ remains in a bounded set of $L^r(Q)$ (r as in Lemma 4).

The proof is straightforward.

PROOF OF THEOREM 1. We can assume that there exists a function ω such that

- (i) ω^ε converges weakly to ω in $L^q(0, T; W_0^{1,q}(\Omega))$ for all $q \in [1, \frac{4}{3}]$.
- (ii) ω^ε converges to ω almost everywhere in Q (Use the compactness' result of [2]).
- (iii) $\operatorname{grad}(L\omega^\varepsilon)$ converges to $\operatorname{grad}(L\omega)$ almost everywhere in Q (use the definition of L , Calderon-Zygmund theorem and the preceding point (ii)).
- (iv) For all φ in $C^\infty(\overline{Q})$, $\omega^\varepsilon B(\omega^\varepsilon, \varphi)$ converges to $\omega B(\omega, \varphi)$ in $L^1(Q)$ -strong.

Multiplying by a suitable function φ (as in Theorem 1) and passing to the limit in (P_ε) , we prove Theorem 1.

REFERENCES

1. B. Michaux and J.M. Rakotoson, (in preparation).
2. R. Temam, *Navier-Stokes Equation*, North-Holland, (1984).
3. G.H. Cottet, Equations de Navier-Stokes dans le plan avec tourbillon initial mesure, *C.R.A.S.* **303**, 105–108 (1986).
4. G.H. Cottet, Two-dimensional incompressible fluid flow with singular initial data, *Technical Report 80*, LMC-IMAG, France, (March 1992).
5. J.M. Rakotoson, A compactness lemma for quasilinear problems: Application to parabolic equations, *J. Funct. Anal.* **106**, 358–374 (1992).